

AGU'98 Special Session
"The Impact of Global Environmental Change on the Hydrological Cycle"

**Freeze/Thaw as a Measure of Global Change Responses of the Boreal
Land Surface Using Spaceborne Radars**

JoBea WAY, Kyle MCDONALD
Jet Propulsion Laboratory, Mail Stop 300-233, 4800 Oak Grove Dr., Pasadena, CA 91109
818-354-8225 and 818-354-3263, 818-354-9476 (FAX), way@jpl.nasa.gov,
kyle.mcdonald@jpl.nasa.gov

Steve FROLKING
Complex Systems Research Center, Morse Hall, University of New Hampshire,
Durham, NH 03824 603-862-0244, 603-862-0188 (FAX), steve.frolking@unh.edu

Steve RUNNING, John KIMBALL
School of Forestry, University of Montana, Missoula, MT 59812-1063 406-243-6311, 406-
243-4510 (FAX), swr@ntsg.umt.edu, johnk@ntsg.umt.edu

Reiner ZIMMERMANN
Institute of Terrestrial Ecology, BITOEK - Plant Ecology II, Dr. Hans Frisch Str. 1,
University of Bayreuth, D-95448 Bayreuth, Germany (49)-921-555624,
reiner.zimmermann@bitoek.uni-bayreuth.de

global change

General Circulation Models (GCMs) project rapid warming in the high latitudes during the next century with major implications on global weather, and carbon and hydrologic cycles. A characteristic of the boreal region that would be impacted by this warming is the temporal and spatial transition of the land surface from a frozen to thawed state. This transition represents the closest analog to a biospheric and hydrologic on/off switch existing in nature, affecting surface meteorological conditions, ecological trace gas dynamics and hydrologic activity profoundly. However, direct evidence to support these boreal warming projections is weak, in large part because of poor climate monitoring capability in these sparsely populated areas. Attempts to overcome the spatial inadequacies of the ground meteorological network using optical/IR satellite systems are severely limited because of poor temporal sampling due to cloud cover and low winter sun angles. In this study we have used microwave data from both spaceborne scatterometers and imaging radars to derive freeze/thaw state. Our work has focused on the boreal regions in Alaska and Canada using 1997/8 NSCAT data and ERS-1/2 data collected over the past 7 years. From these data, landscape freeze/thaw state are being inferred by monitoring shifts in backscatter relative to winter frozen conditions. The inferred landscape freeze/thaw state is validated against temperature measurements obtained from a distributed temperature monitoring network and from meteorological observations located along the spaceborne data transects. A local scale model is used to develop the relationship between freeze/thaw and ecosystem carbon and water flux, and a regional scale model uses the freeze/thaw state as input to estimate regional fluxes. Results from data transects intersecting the Bonanza Creek Experimental Forest in Alaska, and the BOREAS test sites in Canada will be presented.

This work was performed at the Jet Propulsion Laboratory under contract to the National Aeronautics and Space Administration.